

Statement from specialists in nicotine science and public health policy

Dr Margaret Chan
Director General
World Health Organisation
Geneva

CC: FCTC Secretariat, Parties to the FCTC, WHO Regional Offices

26 May 2014

Dear Dr Chan

Reducing the toll of death and disease from tobacco – tobacco harm reduction and the Framework Convention on Tobacco Control (FCTC)

We are writing in advance of important negotiations on tobacco policy later in the year at the FCTC Sixth Conference of the Parties. The work of WHO and the FCTC remains vital in reducing the intolerable toll of cancer, cardiovascular disease and respiratory illnesses caused by tobacco use. As WHO has stated, up to one billion preventable tobacco-related premature deaths are possible in the 21st Century. Such a toll of death, disease and misery demands that we are relentless in our search for all possible practical, ethical and lawful ways to reduce this burden.

It is with concern therefore that a critical strategy appears to have been overlooked or even purposefully marginalised in preparations for FCTC COP-6. We refer to 'tobacco harm reduction' - the idea that the 1.3 billion people who currently smoke could do much less harm to their health if they consumed nicotine in low-risk, non-combustible form.

We have known for years that people 'smoke for the nicotine, but die from the smoke': the vast majority of the death and disease attributable to tobacco arises from inhalation of tar particles and toxic gases drawn into the lungs. There are now rapid developments in nicotine-based products that can effectively substitute for cigarettes but with very low risks. These include for example, e-cigarettes and other vapour products, low-nitrosamine smokeless tobacco such as snus, and other low-risk non-combustible nicotine or tobacco products that may become viable alternatives to smoking in the future. Taken together, these tobacco harm reduction products could play a significant role in meeting the 2025 UN non-communicable disease (NCD) objectives by driving down smoking prevalence and cigarette consumption. Indeed, it is hard to imagine major reductions in tobacco-related NCDs without the contribution of tobacco harm reduction. Even though most of us would prefer people to quit smoking and using nicotine altogether, experience suggests that many smokers cannot or choose not to give up nicotine and will continue to smoke if there is no safer alternative available that is acceptable to them.

We respectfully suggest that the following principles should underpin the public health approach to tobacco harm reduction, with global leadership from WHO:

Statement from specialists in nicotine science and public health policy

1. *Tobacco harm reduction is part of the solution, not part of the problem.* It could make a significant contribution to reducing the global burden of non-communicable diseases caused by smoking, and do so much faster than conventional strategies. If regulators treat low-risk nicotine products as traditional tobacco products and seek to reduce their use without recognising their potential as low-risk alternatives to smoking, they are improperly defining them as part of the problem.
2. *Tobacco harm reduction policies should be evidence-based and proportionate to risk, and give due weight to the significant reductions in risk that are achieved when a smoker switches to a low risk nicotine product.* Regulation should be proportionate and balanced to exploit the considerable health opportunities, while managing residual risks. The architecture of the FCTC is not currently well suited to this purpose.
3. *On a precautionary basis, regulators should avoid support for measures that could have the perverse effect of prolonging cigarette consumption.* Policies that are excessively restrictive or burdensome on lower risk products can have the unintended consequence of protecting cigarettes from competition from less hazardous alternatives, and cause harm as a result. Every policy related to low risk, non-combustible nicotine products should be assessed for this risk.
4. *Targets and indicators for reduction of tobacco consumption should be aligned with the ultimate goal of reducing disease and premature death, not nicotine use per se, and therefore focus primarily on reducing smoking.* In designing targets for the non-communicable disease (NCD) framework or emerging Sustainable Development Goals it would be counterproductive and potentially harmful to include reduction of low-risk nicotine products, such as e-cigarettes, *within these targets*: instead these products should have an important role in *meeting the targets*.
5. *Tobacco harm reduction is strongly consistent with good public health policy and practice and it would be unethical and harmful to inhibit the option to switch to tobacco harm reduction products.* As the WHO's Ottawa Charter states: "*Health promotion is the process of enabling people to increase control over, and to improve, their health*". Tobacco harm reduction allows people to control the risk associated with taking nicotine and to reduce it down to very low or negligible levels.
6. *It is counterproductive to ban the advertising of e-cigarettes and other low risk alternatives to smoking.* The case for banning tobacco advertising rests on the great harm that smoking causes, but no such argument applies to e-cigarettes, for example, which are far more likely to reduce harm by reducing smoking. Controls on advertising to non-smokers, and particularly to young people are certainly justified, but a total ban would have many negative effects, including protection of the cigarette market and implicit support for tobacco companies. It is possible to target advertising at existing smokers where the benefits are potentially huge and the risks minimal. It is inappropriate to apply Article 13 of the FCTC (Tobacco advertising, promotion and sponsorship) to these products.

Statement from specialists in nicotine science and public health policy

7. *It is inappropriate to apply legislation designed to protect bystanders or workers from tobacco smoke to vapour products.* There is no evidence at present of material risk to health from vapour emitted from e-cigarettes. Decisions on whether it is permitted or banned in a particular space should rest with the owners or operators of public spaces, who can take a wide range of factors into account. Article 8 of the FCTC (Protection from exposure to tobacco smoke) should not be applied to these products at this time.
8. *The tax regime for nicotine products should reflect risk and be organised to create incentives for users to switch from smoking to low risk harm reduction products.* Excessive taxation of low risk products relative to combustible tobacco deters smokers from switching and will cause more smoking and harm than there otherwise would be.
9. *WHO and national governments should take a dispassionate view of scientific arguments, and not accept or promote flawed media or activist misinterpretations of data.* For example, much has been made of 'gateway effects', in which use of low-risk products would, it is claimed, lead to use of high-risk smoked products. We are unaware of any credible evidence that supports this conjecture. Indeed, similar arguments have been made about the use of smokeless tobacco in Scandinavia but the evidence is now clear that this product has made a significant contribution to reducing both smoking rates and tobacco-related disease, particularly among males.
10. *WHO and parties to the FCTC need credible objective scientific and policy assessments with an international perspective.* The WHO Study Group on Tobacco Product Regulation (TobReg) produced a series of high quality expert reports between 2005 and 2010. This committee should be constituted with world-class experts and tasked to provide further high-grade independent advice to the WHO and Parties on the issues raised above.

The potential for tobacco harm reduction products to reduce the burden of smoking related disease is very large, and these products could be among the most significant health innovations of the 21st Century – perhaps saving hundreds of millions of lives. The urge to control and suppress them as tobacco products should be resisted and instead regulation that is fit for purpose and designed to realise the potential should be championed by WHO. We are deeply concerned that the classification of these products as tobacco and their inclusion in the FCTC will do more harm than good, and obstruct efforts to meet the targets to reduce non-communicable disease we are all committed to. We hope that under your leadership, the WHO and FCTC will be in the vanguard of science-based, effective and ethical tobacco policy, embracing tobacco harm reduction.

We would be grateful for your considered reaction to these proposals, and we would like to request a meeting with you and relevant staff and a small delegation of signatories to this letter. This statement and any related information will be available on the Nicotine Science and Policy web site (<http://nicotinepolicy.net>) from 29 May 2014.

Yours sincerely,

Statement from specialists in nicotine science and public health policy

Signatories this statement at 26 May 2014

Professor David Abrams

Professor of Health Behavior and Society.
The Johns Hopkins Bloomberg School of
Public Health. Maryland. USA.
Professor of Oncology (adjunct).
Georgetown University Medical Center,
Lombardi Comprehensive Cancer Center.
Washington DC.
United States of America

Professor Tony Axéll

Emeritus Professor Geriatric Dentistry
Consultant in Oral Medicine
Sweden

Professor Pierre Bartsch

Respiratory physician,
Faculty of Medicine
University of Liège
Belgium

Professor Linda Bauld

Professor of Health Policy
Director of the Institute for Social Marketing
Deputy Director, UK Centre for Tobacco
and Alcohol Studies
University of Stirling
United Kingdom

Professor Ron Borland

Nigel Gray Distinguished Fellow in Cancer
Prevention at Cancer Council Victoria
Professorial Fellow School of Population
Health and Department of Information
Systems
University of Melbourne,
Australia

Professor John Britton

Professor of Epidemiology;
Director, UK Centre for Tobacco & Alcohol
Studies,
Faculty of Medicine & Health Sciences
University of Nottingham,
United Kingdom

Associate Professor Chris Bullen

Director, National Institute for Health
Innovation
School of Population Health,
University of Auckland,
New Zealand

Professor Emeritus André Castonguay

Faculty of Pharmacy
Université Laval,
Quebec,
Canada.

Dr Lynne Dawkins

Senior Lecturer in Psychology,
Co-ordinator: Drugs and Addictive
Behaviours Research Group
School of Psychology,
University of East London,
United Kingdom

Professor Ernest Drucker

Professor Emeritus
Department of Family and Social Medicine,
Montefiore Medical Center/Albert Einstein
College of Medicine
Mailman School of Public Health
Columbia University
United States of America

Professor Jean François Etter

Associate Professor
Institut de santé globale,
Faculté de médecine,
Université de Genève,
Switzerland

Dr Karl Fagerström

President, Fagerström Consulting AB,
Vaxholm,
Sweden

Dr Konstantinos Farsalinos

Researcher, Onassis Cardiac Surgery
Center, Athens, Greece
Researcher, University Hospital
Gathuisberg, Leuven,
Belgium

Professor Antoine Flahault

Directeur de l'Institut de Santé Globale
Faculté de Médecine, Université de
Genève, Suisse/ Institute of Global Health,
University of Geneva, Switzerland
Professor of Public Health at the Faculté
de Médecine, Université Paris Descartes,
Sorbonne Paris Cité,
France

Statement from specialists in nicotine science and public health policy

Dr Coral Gartner

Senior Research Fellow
University of Queensland Centre for
Clinical Research
The University of Queensland,
Australia

Dr Guillermo González

Psychiatrist
Comisión de Rehabilitación en Enfermedad
Mental Grave
Clínica San Miguel
Madrid,
Spain

Dr Nigel Gray

Member of Special Advisory Committee on
Tobacco Regulation of the World Health
Organization
Honorary Senior Associate
Cancer Council Victoria
Australia

Professor Peter Hajek

Professor of Clinical Psychology and
Director, Health and Lifestyle Research
Unit
UK Centre for Tobacco and Alcohol
Studies
Wolfson Institute of Preventive Medicine,
Barts and The London School of Medicine
and Dentistry Queen Mary University of
London,
United Kingdom

Professor Wayne Hall

Director and Inaugural Chair, Centre for
Youth Substance Abuse Research
University of Queensland
Australia

Professor John Hughes

Professor of Psychology, Psychiatry and
Family Practice
University of Vermont
United States of America

Professor Martin Jarvis

Emeritus Professor of Health Psychology
Department of Epidemiology & Public
Health
University College London,
United Kingdom

Professor Didier Jayle

Professeur d'addictologie
Conservatoire National des Arts et Métiers
Paris,
France

Dr Martin Juneau

Directeur, Direction de la Prévention
Institut de Cardiologie de Montréal
Professeur Titulaire de Clinique
Faculté de Médecine,
Université de Montréal,
Canada

Dr Michel Kazatchkine

Member of the Global Commission on Drug
Policy
Senior fellow, Global Health Program,
Graduate institute, Geneva,
Switzerland

Professor Demetrios Kouretas

School of Health Sciences and Vice Rector
University of Thessaly,
Greece

Professor Lynn Kozlowski

Dean, School of Public Health and Health
Professions,
Professor of Community Health and Health
Behavior,
University at Buffalo,
State University of New York,
United States of America

Professor Eva Králiková

Institute of Hygiene and Epidemiology
Centre for Tobacco-Dependence
First Faculty of Medicine
Charles University in Prague and General
University Hospital in Prague,
Czech Republic

Professor Michael Kunze

Head of the Institute for Social Medicine
Medical University of Vienna,
Austria

Dr Murray Laugesen

Director
Health New Zealand, Lyttelton,
Christchurch,
New Zealand

Statement from specialists in nicotine science and public health policy

Dr Jacques Le Houezec

Consultant in Public Health, Tobacco dependence, Rennes, France
Honorary Lecturer, UK Centre for Tobacco Control Studies, University of Nottingham, United Kingdom

Dr Kgosi Letlape

President of the Africa Medical Association
Former President of the World Medical Association
Former Chairman of Council of the South African Medical Association
South Africa

Dr Karl Erik Lund

Research director
Norwegian Institute for Alcohol and Drug Research, Oslo, Norway

Dr Gérard Mathern

Président de l'Institut Rhône-Alpes de Tabacologie
Saint-Chamond, France

Professor Richard Mattick

NHMRC Principal Research Fellow
Immediate Past Director NDARC (2001-2009)
National Drug and Alcohol Research Centre (NDARC)
Faculty of Medicine
The University of New South Wales, Australia

Professor Ann McNeill

Professor of Tobacco Addiction
Deputy Director, UK Centre for Tobacco and Alcohol Studies
National Addiction Centre
Institute of Psychiatry
King's College London, United Kingdom

Dr Hayden McRobbie

Reader in Public Health Interventions, Wolfson Institute of Preventive Medicine, Queen Mary University of London, United Kingdom

Dr Anders Milton

Former President of the Swedish Red Cross
Former President and Secretary of the Swedish Medical Association
Former Chairman of the World Medical Association
Owner & Principal Milton Consulting, Sweden

Professor Marcus Munafò

Professor of Biological Psychology
MRC Integrative Epidemiology Unit at the University of Bristol
UK Centre for Tobacco and Alcohol Studies
School of Experimental Psychology
University of Bristol, United Kingdom

Professor David Nutt

Chair of the Independent Scientific Committee on Drugs (UK)
Edmund J Safra Professor of Neuropsychopharmacology
Head of the Department of Neuropsychopharmacology and Molecular Imaging
Imperial College London, United Kingdom

Dr Gaston Ostiguy

Professeur agrégé
Directeur de la Clinique de cessation tabagique
Centre universitaire de santé McGill (CUSM)
Institut thoracique de Montréal, Canada

Professor Riccardo Polosa

Director of the Institute for Internal Medicine and Clinical Immunology, University of Catania, Italy.

Dr Lars Ramström

Director
Institute for Tobacco Studies
Täby, Sweden

Statement from specialists in nicotine science and public health policy

Dr Martin Raw

Special Lecturer
UK Centre for Tobacco and Alcohol
Studies
Division of Epidemiology and Public Health
University of Nottingham,
United Kingdom

Professor Andrzej Sobczak

Department of General and Inorganic
Chemistry,
Faculty of Pharmacy and Laboratory
Medicine,
Medical University of Silesia, Katowice,
Poland
Institute of Occupational Medicine and
Environmental Health
Sosnowiec,
Poland

Professor Gerry Stimson

Emeritus Professor, Imperial College
London;
Visiting Professor, London School of
Hygiene and Tropical Medicine
United Kingdom

Professor Tim Stockwell

Director, Centre for Addictions Research of
BC
Professor, Department of Psychology
University of Victoria, British Columbia,
Canada

Professor David Sweanor

Adjunct Professor, Faculty of Law,
University of Ottawa
Special Lecturer, Division of Epidemiology
and Public Health,
University of Nottingham,
United Kingdom

Professor Umberto Tirelli

Director Department of Medical Oncology
National Cancer Institute of Aviano
Italy

Professor Umberto Veronesi

Scientific Director
IEO Istituto Europeo di Oncologia
Former Minister of Health,
Italy

Professor Kenneth Warner

Avedis Donabedian Distinguished
University Professor of Public Health
Professor, Health Management & Policy
School of Public Health
University of Michigan
United States of America

Professor Robert West

Professor of Health Psychology and
Director of Tobacco Studies
Health Behaviour Research Centre,
Department of Epidemiology & Public
Health,
University College London
United Kingdom

Professor Dan Xiao

Director of Department Epidemiology
WHO Collaborating Center for Tobacco or
Health
Beijing Institute of Respiratory Medicine,
Beijing Chao-Yang Hospital,
China

Dr Derek Yach

Former Executive Director, Non-
Communicable Diseases
Former Head of Tobacco Free Initiative,
World Health Organisation (1995-2004)
Senior Vice President Vitality Group plc
Director, Vitality Institute for Health
Promotion
United States of America